

C_k -MOVES ON SPATIAL THETA-CURVES AND VASSILIEV INVARIANTS

AKIRA YASUHARA

Department of Mathematics, Tokyo Gakugei University
 Nukuikita 4-1-1, Koganei, Tokyo 184-8501, Japan

Current address, October 1, 1999 to September 30, 2001:
 Department of Mathematics, The George Washington University
 Washington, DC 20052, USA
 e-mail: yasuhara@u-gakugei.ac.jp

ABSTRACT

The C_k -equivalence is an equivalence relation generated by C_k -moves defined by Habiro. Habiro showed that the set of C_k -equivalence classes of the knots forms an abelian group under the connected sum and it can be classified by the additive Vassiliev invariant of order $\leq k-1$. We see that the set of C_k -equivalence classes of the spatial θ -curves forms a group under the vertex connected sum and that if the group is abelian, then it can be classified by the additive Vassiliev invariant of order $\leq k-1$. However the group is not necessarily abelian. In fact, we show that it is nonabelian for $k \geq 12$. As an easy consequence, we have the set of C_k -equivalence classes of m -string links, which forms a group under the composition, is nonabelian for $k \geq 12$ and $m \geq 2$.

1. C_k -moves and Vassiliev invariants of spatial θ -curves

A *tangle* T is a disjoint union of properly embedded arcs in the unit 3-ball B^3 . A *local move* is a pair of tangles (T_1, T_2) with $\partial T_1 = \partial T_2$ such that for each component t of T_1 there exists a component u of T_2 with $\partial t = \partial u$. Two local moves (T_1, T_2) and (U_1, U_2) are *equivalent* if there is an orientation preserving self-homeomorphism $\psi : B^3 \rightarrow B^3$ such that $\psi(T_i)$ and U_i are ambient isotopic in B^3 relative to ∂B^3 for $i = 1, 2$. Here $\psi(T_i)$ and U_i are *ambient isotopic in B^3 relative to ∂B^3* if $\psi(T_i)$ is deformed to U_i by an ambient isotopy of B^3 that is pointwisely fixed on ∂B^3 .

Let (T_1, T_2) be a local move, t_1 a component of T_1 and t_2 a component of T_2 with $\partial t_1 = \partial t_2$. Let N_1 and N_2 be regular neighbourhoods of t_1 and t_2 in $(B^3 - T_1) \cup t_1$ and $(B^3 - T_2) \cup t_2$ respectively such that $N_1 \cap \partial B^3 = N_2 \cap \partial B^3$. Let α be a disjoint union of properly embedded arcs in $B^2 \times [0, 1]$ as illustrated in Fig. 1.1. Let $\psi_i : B^2 \times [0, 1] \rightarrow N_i$ be a homeomorphism with $\psi_i(B^2 \times \{0, 1\}) = N_i \cap \partial B^3$ for $i = 1, 2$. Suppose that

2000 *Mathematics Subject Classification*: Primary 57M25; Secondary 57M27

Keywords and Phrases: spatial theta-curve, C_n -move, Vassiliev invariant, finite type invariant

$\psi_1(\partial\alpha) = \psi_2(\partial\alpha)$ and $\psi_1(\alpha)$ and $\psi_2(\alpha)$ are ambient isotopic in B^3 relative to ∂B^3 . Then we say that a local move $((T_1 - t_1) \cup \psi_1(\alpha), (T_2 - t_2) \cup \psi_2(\alpha))$ is a *double* of (T_1, T_2) with respect to the components t_1 and t_2 .




Fig. 1.1

A C_1 -move is a local move (T_1, T_2) as illustrated in Fig. 1.2. A double of a C_k -move is called a C_{k+1} -move. Note that, for each natural number k , there are only finitely many C_k -moves up to equivalence. It is easy to see that if (T_1, T_2) is a C_n -move, then (T_2, T_1) is equivalent to a C_n -move (but possibly not equivalent to itself). The definition of C_k -move follows that in [2], and is different from the one in [3]. However by an easy induction on k it is shown that these two definitions are essentially same. In [3], a C_k -move is called a *simple* C_k -move, and a C_k -move means a *parallel* of a C_k -move. The definition of parallel of a local move appears in Section 3.




Fig. 1.2

Let G be a graph with labeled vertices and edges. Let $f : G \rightarrow S^3$ be an embedding of G into the oriented three sphere S^3 . The embedding f is called a *spatial graph*.

Let f_1 and f_2 be spatial graphs. We say that f_2 is *obtained from f_1 by a local move (T_1, T_2)* if there is an orientation preserving embedding $h : B^3 \rightarrow S^3$ such that $f_i(G) \cap h(B^3) = h(T_i)$ for $i = 1, 2$ and $f_1(G - h(B^3)) = f_2(G - h(B^3))$ together with the labels of vertices and edges. Two spatial graphs f_1 and f_2 are C_k -equivalent if f_2 is obtained from f_1 by a finite sequence of C_k -moves and ambient isotopies. We note that the relation is an equivalence relation on spatial graphs. For a spatial graph f , let $[f]_k$ denote the C_k -equivalence class that contains f . It is known that C_k -equivalence implies C_{k-1} -equivalence [3], [16].

Let l be a positive integer and $k_1, \dots, k_l$ positive integers. Suppose that for each $P \subset \{1, \dots, l\}$, a spatial graph f_P in S^3 is assigned. Suppose that there are mutually disjoint, orientation preserving embeddings $h_i : B^3 \rightarrow S^3$ ($i = 1, \dots, l$) such that

- (1) $f_\emptyset(G) - \bigcup_{i=1}^l h_i(B^3) = f_P(G) - \bigcup_{i=1}^l h_i(B^3)$ together with the labels for any subset

$P \subset \{1, \dots, l\}$,

(2) $(h_i^{-1}(f_\emptyset(G)), h_i^{-1}(f_{\{1, \dots, l\}}(G)))$ is a C_{k_i} -move ($i = 1, \dots, l$), and

(3) $f_P(G) \cap h_i(B^3) = \begin{cases} f_{\{1, \dots, l\}}(G) \cap h_i(B^3) & \text{if } i \in P, \\ f_\emptyset(G) \cap h_i(B^3) & \text{otherwise.} \end{cases}$

Then we call the set $\{f_P | P \subset \{1, \dots, l\}\}$ a *singular spatial graph of type $(k_1, \dots, k_l)$* .

The θ -curve is a graph θ with two vertices v_1, v_2 and three edges e_1, e_2, e_3 each of which joins v_1 and v_2 . When $G = \theta$, a spatial graph and singular spatial graph are called *spatial θ -curve* and *singular spatial θ -curve* respectively. A spatial θ -curve is *trivial* if its image is contained in a 2-sphere in S^3 . In the remainder of this section, we consider only the case that a graph is the θ -curve.

Let Γ be the set of all spatial θ -curve types in S^3 and $\mathbb{Z}\Gamma$ the free abelian group generated by Γ . For a singular spatial θ -curve $\mathbf{f} = \{f_P | P \subset \{1, \dots, l\}\}$ of type $(k_1, \dots, k_l)$, we define an element $\kappa(\mathbf{f})$ of $\mathbb{Z}\Gamma$ by

$$\kappa(\mathbf{f}) = \sum_{P \subset \{1, \dots, l\}} (-1)^{|P|} f_P.$$

Let $\mathcal{V}(k_1, \dots, k_l)$ be the subgroup of $\mathbb{Z}\Gamma$ generated by all $\kappa(\mathbf{f})$ where $\mathbf{f}$ varies over all singular spatial θ -curves of type $(k_1, \dots, k_l)$.

For two spatial θ -curves f_1 and f_2 , remove small balls centred at $f_1(v_2)$ and $f_2(v_1)$ from S^3 , then identify the boundaries so that the images of i -th edge are joined for each i . Then we obtain a new spatial θ -curve. We call this embedding the *vertex connected sum* of f_1 and f_2 , and denote by $f_1 \# f_2$. The vertex connected sum is well-defined up to ambient isotopy [18]. Let $f_1 \# f_2$ be the vertex connected sum of two spatial θ -curves f_1 and f_2 . Then $f_1 \# f_2 - f_1 - f_2 \in \mathbb{Z}\Gamma$ is called a *composite relator*. Let $\mathcal{R}_\#$ be the subgroup of $\mathbb{Z}\Gamma$ generated by all composite relators.

Let $\iota : \Gamma \rightarrow \mathbb{Z}\Gamma$ be the natural inclusion map. Let $\pi : \mathbb{Z}\Gamma \rightarrow \mathbb{Z}\Gamma / \mathcal{V}(k_1, \dots, k_l)$ and $\lambda : \mathbb{Z}\Gamma / \mathcal{V}(k_1, \dots, k_l) \rightarrow \mathbb{Z}\Gamma / (\mathcal{V}(k_1, \dots, k_l) + \mathcal{R}_\#)$ be the quotient homomorphisms. Then the composite maps $\pi \circ \iota : \Gamma \rightarrow \mathbb{Z}\Gamma / \mathcal{V}(k_1, \dots, k_l)$ and $\lambda \circ \pi \circ \iota : \Gamma \rightarrow \mathbb{Z}\Gamma / (\mathcal{V}(k_1, \dots, k_l) + \mathcal{R}_\#)$ are called the *universal Vassiliev invariant of type $(k_1, \dots, k_l)$* and *universal additive Vassiliev invariant of type $(k_1, \dots, k_l)$* respectively. We denote them by $v_{(k_1, \dots, k_l)}$ and $w_{(k_1, \dots, k_l)}$ respectively. In the case of knots, these are same invariants as defined by K. Taniyama and the author [17]. Similarly, we can also define $\mathcal{V}(k_1, \dots, k_l)$ and the universal Vassiliev invariant $v_{(k_1, \dots, k_l)}$ for the embeddings of any graph. Since a C_1 -move is a crossing change we see that a singular spatial graph of type $(\underbrace{1, \dots, 1}_l)$ is essentially the same as a

singular spatial graph with l crossing vertices in the sense of T. Stanford [12]. Therefore we see that $v_{(1, \dots, 1)}$ is the universal Vassiliev invariant of order $\leq l - 1$. Note that $v_{(1, \dots, 1)}(f_1) = v_{(1, \dots, 1)}(f_2)$ if and only if $v(f_1) = v(f_2)$ for any Vassiliev invariant v of order $\leq l - 1$. In the case of links, $v_{(2, \dots, 2)}$ is the same as that defined in [7], [13]. In [16] Taniyama and the author defined finite type invariants of order $(k; n)$ for the embeddings of a graph, which are essentially same as $v_{(\underbrace{n-1, \dots, n-1}_{k+1})}$.

By the arguments similar to that in Proofs of Theorems 1.1 and 1.2 in [17] and that in Proof of Theorem 1.4 in [17], we have the following two theorems.

Theorem 1.1. *Let $k_1, \dots, k_l$ be positive integers and $k = k_1 + \dots + k_l$. Then the followings hold.*

- (1) $\mathcal{V}(k) \subset \mathcal{V}(k_1, \dots, k_l) \subset \mathcal{V}(\underbrace{1, \dots, 1}_k)$.
- (2) $\mathcal{V}(k_1, \dots, k_l) + \mathcal{R}_\# = \mathcal{V}(k) + \mathcal{R}_\#$. $\square$

Remark. Theorem 1.1(1) holds for the spatial embeddings of any graph.

Theorem 1.2. *The C_k -equivalence classes of the spatial θ -curves forms a group with the unit element $[f_0]_k$ under the vertex connected sum, where f_0 is a trivial θ -curve.* $\square$

We denote by G_k this group. Let $\varphi : G_k \rightarrow \mathbb{Z}\Gamma/\mathcal{V}(k)$ be a map induced by the inclusion $\iota : \Gamma \rightarrow \mathbb{Z}\Gamma$. By Theorem 1.2, φ is a well-defined, epimorphism. (In fact, φ is an isomorphism, see the remark after Corollary 1.4.) Since $\varphi([G_k, G_k]) = \mathcal{R}_\#$, by Theorem 1.1 (2), we have the following theorem.

Theorem 1.3. *Let $k_1, \dots, k_l$ be positive integers and $k = k_1 + \dots + k_l$. Then $G_k/[G_k, G_k]$ is isomorphic to $\mathbb{Z}\Gamma/(\mathcal{V}(k_1, \dots, k_l) + \mathcal{R}_\#)$.* $\square$

Let f_1 and f_2 be spatial θ -curves and $k = k_1 + \dots + k_l$. If f_1 and f_2 are C_k -equivalent, then by Theorem 1.1 (1), $f_1 - f_2 \in \mathcal{V}(k) \subset \mathcal{V}(k_1, \dots, k_l)$. Therefore we have $v_{(k_1, \dots, k_l)}(f_1) = v_{(k_1, \dots, k_l)}(f_2)$. On the other hand, if $v_{(k_1, \dots, k_l)}(f_1) = v_{(k_1, \dots, k_l)}(f_2)$, then $w_{(k_1, \dots, k_l)}(f_1) = w_{(k_1, \dots, k_l)}(f_2)$. Hence we have $f_1 - f_2 \in \mathcal{V}(k_1, \dots, k_l) + \mathcal{R}_\#$. If G_k is abelian group, i.e., $[G_k, G_k] = \{id\}$, then by Theorem 1.3, f_1 and f_2 are C_k -equivalent. So we have the following corollary.

Corollary 1.4. *Let $k_1, \dots, k_l$ be positive integers and $k = k_1 + \dots + k_l$. Let f_1 and f_2 be spatial θ -curves. If G_k is an abelian group, then the following conditions are mutually equivalent.*

- (1) f_1 and f_2 are C_k -equivalent,
- (2) $v_{(k_1, \dots, k_l)}(f_1) = v_{(k_1, \dots, k_l)}(f_2)$,
- (3) $w_{(k_1, \dots, k_l)}(f_1) = w_{(k_1, \dots, k_l)}(f_2)$. $\square$

Remark. Let f_1 and f_2 be spatial graphs (not necessarily θ -curve). If $f_1 - f_2 \in \mathcal{V}(k)$, then there are singular spatial graphs $\mathbf{f}^i$'s of type (k) and integers x_i 's such that $f_1 - f_2 = \sum x_i \kappa(\mathbf{f}^i)$. By induction on $\sum |x_i|$, we see that f_1 and f_2 are C_k -equivalent. Since $f_1 - f_2 \in \mathcal{V}(k)$ if f_1 and f_2 are C_k -equivalent, we have the following: Two spatial graphs f_1 and f_2 are C_k -equivalent if and only if $v_{(k)}(f_1) = v_{(k)}(f_2)$.

Theorem 1.5. *Let f_0 be a trivial spatial θ -curve. Then the followings hold.*

- (1) For each $f \in [f_0]_k$, $[f]_{k+2}$ belongs to the center of G_{k+2} .
- (2) If $2l \geq k$, then the set $H_k^l = \{[f]_k | f \in [f_0]_l\}$ is an abelian subgroup of G_k .

By [15] and [8], we have $[f_0]_1 = [f_0]_2 = \Gamma$. Hence, by Theorem 1.5(2), $H_4^2 = G_4$ is abelian. By Corollary 1.4, we have

Corollary 1.6. *Let $k_1, \dots, k_l$ ($l \leq 4$) be positive integers and $k = k_1 + \dots + k_l \leq 4$. Let f_1 and f_2 be spatial θ -curves. Then the following conditions are mutually equivalent.*

- (1) f_1 and f_2 are C_k -equivalent,
- (2) $v_{(k_1, \dots, k_l)}(f_1) = v_{(k_1, \dots, k_l)}(f_2)$,
- (3) $w_{(k_1, \dots, k_l)}(f_1) = w_{(k_1, \dots, k_l)}(f_2)$. $\square$

Remark. As a special case of Corollary 1.6, we see that for $k \leq 4$ two spatial θ -curves are C_k -equivalent if and only if the universal (additive) Vassiliev invariant of order $\leq k-1$ are equal. Meanwhile, a basis for the space of Vassiliev invariants of order ≤ 4 is known [4],[6].

Theorem 1.7. *Let f_0 be a trivial spatial θ -curve, and let f_1 and f_2 be in $[f_0]_k$. Then f_1 and f_2 are C_{2k} -equivalent if and only if $v_{(k,k)}(f_1) = v_{(k,k)}(f_2)$.*

As we saw before, G_k is abelian for $k \geq 4$. However G_k is not necessarily abelian. In fact, we have the following theorem.

Theorem 1.8. *The group G_k is nonabelian for any $k \geq 12$.*

Remarks (1) If G_k is abelian, then so is $G_{k'}$ for any $k'(< k)$.

(2) In the proof of Theorem 1.8, we see that there are two spatial θ -curves g and h such that $g - h$ is in $\mathcal{R}_\#$ and is not in $\mathcal{V}(\underbrace{1, \dots, 1}_k)$ for $k \geq 12$. Hence $v_{(k_1, \dots, k_l)}(g) \neq v_{(k_1, \dots, k_l)}(h)$

for $k_1 + \dots + k_l \geq 12$ by Theorem 1.1(1), while $w_{(k_1, \dots, k_l)}(g) = w_{(k_1, \dots, k_l)}(h)$ for any $k_1, \dots, k_l$. In contrast, for any knots K and K' , $v_{(k_1, \dots, k_l)}(K) = v_{(k_1, \dots, k_l)}(K')$ if and only if $w_{(k_1, \dots, k_l)}(K) = w_{(k_1, \dots, k_l)}(K')$ [17].

(3) Habiro showed the set of C_k -equivalence classes $S_k(m)$ of m -string links forms a group under the composition [3]. By considering the complement of a regular neighbourhood of one of edges, we have that there is a surjection from the 2-string links to the spatial θ -curves. Since the surjection induces an epimorphism from $S_k(2)$ to G_k and since there is an epimorphism from $S_k(m)$ ($m > 2$) to $S_k(2)$, $S_k(m)$ is nonabelian for any $m \geq 2$ and $k \geq 12$.

The following is still open.

Problem. Find the minimum number k ($5 \leq k \leq 12$) such that G_k is nonabelian.

2. Band description of spatial graphs

A C_1 -link model is a pair (α, β) where α is a disjoint union of properly embedded arcs in B^3 and β is a disjoint union of arcs on ∂B^3 with $\partial\alpha = \partial\beta$ as illustrated in Fig. 2.1. Suppose that a C_k -link model (α, β) is defined where α is a disjoint union of $k+1$ properly embedded arcs in B^3 and β is a disjoint union of $k+1$ arcs on ∂B^3 with $\partial\alpha = \partial\beta$ such that $\alpha \cup \beta$ is a disjoint union of $k+1$ circles. Let γ be a component of $\alpha \cup \beta$ and W a regular neighbourhood of γ in $(B^3 - (\alpha \cup \beta)) \cup \gamma$. Let V be an oriented solid torus, D a disk in ∂V , α_0 properly embedded arcs in V and β_0 arcs on D as illustrated in Fig. 2.2. Let $\psi : V \rightarrow W$ be an orientation preserving homeomorphism such that

$\psi(D) = W \cap \partial B^3$ and $\psi(\alpha_0 \cup \beta_0)$ bounds disjoint disks in B^3 . Then we call the pair $((\alpha - \gamma) \cup \psi(\alpha_0), (\beta - \gamma) \cup \psi(\beta_0))$ a C_{k+1} -link model. A link model is a C_k -link model for some k . It is known that, for a C_k -link model (α, β) , the local move $(\alpha, \hat{\beta})$ is equivalent to a C_k -move [17], where $\hat{\beta}$ is a slight push in of β .




Fig. 2.1




Fig. 2.2

Let f_1 be a spatial θ -curve, and let $(\alpha_1, \beta_1), \dots, (\alpha_l, \beta_l)$ be link models. Let $\psi_i : B^3 \rightarrow S^3$ ($i = 1, \dots, l$) be mutually disjoint, orientation preserving embeddings, and let $b_{1,1}, b_{1,2}, \dots, b_{1,\rho(1)}, b_{2,1}, b_{2,2}, \dots, b_{2,\rho(2)}, \dots, b_{l,1}, b_{l,2}, \dots, b_{l,\rho(l)}$ be mutually disjoint disks embedded in S^3 . Suppose that they satisfy the following conditions;

- (1) $\psi_i(B^3) \cap f_1(\theta) = \emptyset$ for each i ,
- (2) $b_{i,k} \cap f_1(\theta) = \partial b_{i,k} \cap (f_1(\theta) - f(v_1 \cup v_2))$ is an arc for each i, k ,
- (3) $b_{i,k} \cap (\bigcup_{j=1}^l \psi_j(B^3)) = \partial b_{i,k} \cap \psi_i(B^3)$ is a component of $\psi_i(\beta_i)$ for each i, k ,
- (4) $(\bigcup_{k=1}^{\rho(i)} b_{i,k}) \cap \psi_i(B^3) = \psi_i(\beta_i)$ for each i .

Let f_2 be a spatial θ -curve defined by

$$f_2(\theta) = f_1(\theta) \cup \left(\bigcup_{i,k} \partial b_{i,k} \right) \cup \left(\bigcup_{i=1}^l \psi_i(\alpha_i) \right) - \bigcup_{i,k} \text{int}(\partial b_{i,k} \cap f_1(\theta)) - \bigcup_{i=1}^l \psi_i(\text{int} \beta_i),$$

where the labels of $f_2(\theta)$ coincides that of $f_1(\theta)$ on $f_1(\theta) - \bigcup_{i,k} b_{i,k}$. When (α_i, β_i) is a C_k -link model, we call $\psi_i(B^3)$ a C_k -link ball. We set $\mathcal{B}_i = ((\alpha_i, \beta_i), \psi_i, \{b_{i,1}, \dots, b_{i,\rho(i)}\})$ and call $\mathcal{B}_i$ a C_k -chord when (α_i, β_i) is a C_k -link model. We denote f_2 by $\Omega(f_1; \{\mathcal{B}_1, \dots, \mathcal{B}_l\})$ and call it a *band description* of f_2 . We also say f_2 is a *band sum* of f_1 and link models $(\alpha_1, \beta_1), \dots, (\alpha_l, \beta_l)$.

By the arguments similar to that in Proof of Lemma 3.6 [17], we have

Lemma 2.1. *Two spatial θ -curves f_1 and f_2 are C_k -equivalence if and only if there are spatial θ -curves f'_1 and f'_2 such that f'_i is ambient isotopic to f_i ($i = 1, 2$) and f'_2 is a band sum of f'_1 and some C_k -link models. $\square$*

In the following lemma, the former assertion follows directly from Lemma 3.9 in [17] and the latter can be shown by the similar arguments as in proof of Lemma 3.9 in [17].

Lemma 2.2. *A local move as illustrated in Fig. 2.3 (resp. Fig. 2.4) is realized by a C_{j+k} -move (resp. C_{j+k+1} -move). $\square$*




Fig. 2.3




Fig. 2.4

Proof of Theorem 1.5. (1) Suppose $f \in [f_0]_k$, then by Lemma 2.1, we may assume that f is a band sum of f_0 and some C_k -link models. Let g be a spatial θ -curve. Since $\Gamma = [f_0]_2$, we may suppose g is a band sum of f_0 and some C_2 -link models. It is not hard to see that $f \# g$ and $g \# f$ are transposed each other by the moves as in Figs. 2.3 and 2.4, where we consider the case $j = 2$, and ambient isotopies. Thus by Lemma 2.2 we have $f \# g$ is C_{k+2} -equivalent to $g \# f$. Hence we have $[f]_{k+2}[g]_{k+2} = [f \# g]_{k+2} = [g \# f]_{k+2} = [g]_{k+2}[f]_{k+2}$.

(2) Suppose that both $[f_1]_k$ and $[f_2]_k$ belong to H_k^l . Then we note that $[f_1]_k[f_2]_k = [f_1 \# f_2]_k \in H_k^l$. If $[f]_k$ belongs to H_k^l , then by Lemma 2.1, we may assume that $f = \Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\})$ and $f_0 = \Omega(f; \{\mathcal{B}'_1, \dots, \mathcal{B}'_m\})$ for some C_l -chords $\mathcal{B}_1, \dots, \mathcal{B}_n, \mathcal{B}'_1, \dots, \mathcal{B}'_m$. By using Sublemma 3.5 in [17] repeatedly, there are C_l -chords $\mathcal{B}''_1, \dots, \mathcal{B}''_m$ such that f_0 is ambient isotopic to $\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n, \mathcal{B}''_1, \dots, \mathcal{B}''_m\})$. By Lemma 2.2, we can deform $\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n, \mathcal{B}''_1, \dots, \mathcal{B}''_m\})$ into $\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\}) \# \Omega(f_0; \{\mathcal{B}''_1, \dots, \mathcal{B}''_m\})$ by C_{2l} -moves and ambient isotopies, i.e., f_0 and $f \# \Omega(f_0; \{\mathcal{B}''_1, \dots, \mathcal{B}''_m\})$ are C_{2l} -equivalent. Since $2l \geq k$, f_0 and $f \# \Omega(f_0; \{\mathcal{B}''_1, \dots, \mathcal{B}''_m\})$ are C_k -equivalent. This implies that $[\Omega(f_0; \{\mathcal{B}''_1, \dots, \mathcal{B}''_m\})]_k = [f]_k^{-1} \in H_k^l$. Therefore H_k^l is a subgroup of G_k . By the arguments similar to that in (1), we see that H_k^l is abelian. $\square$

Let v be an invariant of the embeddings of a graph that takes values in an abelian group. We call v a *Vassiliev invariant of type* $(k_1, \dots, k_l)$ if, for any singular spatial graph $\{f_P | P \subset \{1, \dots, l\}\}$ of type $(k_1, \dots, k_l)$,

$$\sum_{P \subset \{1, \dots, l\}} (-1)^{|P|} v(f_P) = 0.$$

Proof of Theorem 1.7. The ‘only if’ part follows from Theorem 1.1(1). We shall show ‘if’ part. Let $\varphi : \Gamma \rightarrow H_{2k}^k$ be a map defined as follows:

$$\varphi(f) = \begin{cases} [f]_{2k} & \text{if } f \in [f_0]_k, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly, φ is an invariant. Now we will show that φ is a Vassiliev invariant of type (k, k) . Let $\{h_P | P \subset \{1, 2\}\}$ be a singular spatial θ -curve of type (k, k) . Since $\{h_P | P \subset \{1, 2\}\} \subset [h_\emptyset]_k$, we have $\varphi(h_P) = 0$ if $[h_\emptyset]_k \neq [f_0]_k$. So we may suppose that $[h_\emptyset]_k = [f_0]_k$. Then we have

$$\sum_{P \subset \{1, 2\}} (-1)^{|P|} \varphi(h_P) = [h_\emptyset]_{2k} - [h_{\{1\}}]_{2k} - [h_{\{2\}}]_{2k} + [h_{\{1, 2\}}]_{2k}.$$

Since $[h_\emptyset]_k = [f_0]_k$, by Lemma 2.1, we may assume that $h_\emptyset = \Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\})$ for some C_k -chords $\mathcal{B}_1, \dots, \mathcal{B}_n$. By Sublemma 3.1 in [17] (or Lemma 3.7 in the next section), there are C_k -chords $\mathcal{B}'_1, \mathcal{B}'_2$ such that $\Omega(h_\emptyset; \cup_{i \in P} \{\mathcal{B}'_i\})$ is ambient isotopic to h_P . By the arguments similar to that in the proof of Sublemma 3.5 in [17], we see that there are C_k -chords $\mathcal{B}''_1, \mathcal{B}''_2$ such that $\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\} \cup (\cup_{i \in P} \{\mathcal{B}''_i\}))$ is ambient isotopic to h_P . Since $\mathcal{B}_1, \dots, \mathcal{B}_n, \mathcal{B}''_1, \mathcal{B}''_2$ are C_k -chords, by Lemma 2.2, we have

$$[\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\} \cup \{\mathcal{B}''_1, \mathcal{B}''_2\})]_{2k} = [\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\}) \# \Omega(f_0; \{\mathcal{B}''_1\}) \# \Omega(f_0; \{\mathcal{B}''_2\})]_{2k},$$

and

$$[\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\} \cup \{\mathcal{B}''_i\})]_{2k} = [\Omega(f_0; \{\mathcal{B}_1, \dots, \mathcal{B}_n\}) \# \Omega(f_0; \{\mathcal{B}''_i\})]_{2k} \quad (i = 1, 2).$$

Hence we have

$$\begin{aligned} \sum_{P \subset \{1, 2\}} (-1)^{|P|} \varphi(h_P) &= [f_\emptyset]_{2k} - [f_\emptyset \# \Omega(f_0; \{\mathcal{B}''_1\})]_{2k} - [f_\emptyset \# \Omega(f_0; \{\mathcal{B}''_2\})]_{2k} \\ &\quad + [f_\emptyset \# \Omega(f_0; \{\mathcal{B}''_1\}) \# \Omega(f_0; \{\mathcal{B}''_2\})]_{2k} = 0 \in H_{2k}^k. \end{aligned}$$

Therefore φ is a Vassiliev invariant of type (k, k) . This and the assumption $v_{(k, k)}(f_1) = v_{(k, k)}(f_2)$ imply $\varphi(f_1) = \varphi(f_2)$. By the definition of φ , we have $[f_1]_{2k} = [f_2]_{2k}$. $\square$

3. Disk/band surfaces and Vassiliev invariants of spatial graphs

A graph G is *trivalent* if the valence of any vertex of G is equal to 3. A graph G is *planar* if there exists an embedding $f_0 : G \rightarrow \mathbb{R}^2$. A connected, planar graph G is said to be *prime* if, for any embedding $f_0 : G \rightarrow \mathbb{R}^2$, there exist no simple closed curves C in $\mathbb{R}^2$ satisfying either the following (1) or (2) (cf. [14], [5]), where A, B are the two components of $\mathbb{R}^2 - C$.

- (1) C meets $f_0(G)$ in a single point such that both $A \cap f_0(G)$ and $B \cap f_0(G)$ are non-empty.
- (2) C meets $f_0(G)$ in two points such that both $A \cap f_0(G)$, $B \cap f_0(G)$ are neither empty nor single open arcs.

For any connected, planar graph G , we fix a planar embedding $f_0 : G \rightarrow \mathbb{R}^2$ arbitrarily. The image $f_0(G)$ has complementary domains $D_1, D_2, \dots, D_n$ that are bounded and one unbounded D_0 . The preimage $c_i = f_0^{-1}(\partial D_i)$ is a 1-complex which can be viewed as a 1-cycle in $H_1(G; \mathbb{Z})$. We call c_i ($i \neq 0$), c_0 respectively a *boundary cycle* and the *outermost cycle* in G with respect to f_0 .

For a spatial embedding $f : G \rightarrow S^3$ of a graph G , a *disk/band surface* S of $f(G)$ is a compact, orientable surface in S^3 such that $f(G)$ is a deformation retract of S contained in $\text{int} S$ [5].

In [10], T. Soma, H. Sugai and the author showed the following theorem.

Theorem 3.1. ([10, Theorem 1]) *Suppose that G is a connected, planar, prime and trivalent graph, and $f_0 : G \rightarrow \mathbb{R}^2$ is an embedding. Then, for any embedding $f : G \rightarrow S^3$, there exists the unique disk/band surface S of $f(G)$ up to ambient isotopy of which the Seifert pairings satisfying the following equation.*

$$\langle f(c_i), f(c_j) \rangle_S = \begin{cases} -\text{lk}(f(c_i), f(c_0)) - \sum_{c_i \cap c_k = \emptyset} \text{lk}(f(c_i), f(c_k)) & \text{if } i = j \text{ and } c_i \cap c_0 = \emptyset \\ 0 & \text{if } i \neq j \text{ and } c_i \cap c_j \neq \emptyset \\ 0 & \text{if } i = j \text{ and } c_i \cap c_0 \neq \emptyset \\ \text{lk}(f(c_i), f(c_j)) & \text{if } c_i \cap c_j = \emptyset, \end{cases}$$

where c_i, c_j, c_k are boundary cycles and c_0 is the outermost cycle with respect to f_0 . $\square$

We call the disk/band surface above the *canonical disk/band surface* for f . Note that the Seifert linking form of the canonical disk/band surface depends only on the linking numbers of pairs of disjoint cycles. If G is the θ -curve or the complete graph with 4 vertices, then the canonical disk/band surface is same as the *disk/band surface with zero Seifert linking form* that is defined in [5]. By the proof of Theorem 1 in [10], we note that the canonical disk/band surface is given as the image of an embedding of the regular neighborhood S_0 of $f_0(G)$ in $\mathbb{R}^2$. Thus by fixing orientation and label of ∂S_0 , we have an ordered, oriented link as the image of an embedding of ∂S_0 . From now on, we always assume that, for each graph G , ∂S_0 has fixed orientation and label.

Let (T_1, T_2) be a local move, $t_{11}, \dots, t_{1n}$ the components of T_1 and $t_{21}, \dots, t_{2n}$ the components of T_2 with $\partial t_{1i} = \partial t_{2i}$ ($i = 1, \dots, n$). Let N_{1i} and N_{2i} be regular neighbourhoods of t_{1i} and t_{2i} in B^3 respectively such that $N_{1i} \cap \partial B^3 = N_{2i} \cap \partial B^3$ ($i = 1, \dots, n$) and $N_{1i} \cap N_{1j} = N_{2i} \cap N_{2j} = \emptyset$ ($1 \leq i < j \leq n$). Let α_i ($i = 1, \dots, n$) be disjoint union of properly embedded l_i arcs in $B^2 \times [0, 1]$ as illustrated in Fig. 3.1. Let $\psi_{ji} : B^2 \times [0, 1] \rightarrow N_{ji}$ be a homeomorphism with $\psi_{ji}(B^2 \times \{0, 1\}) = N_{ji} \cap \partial B^3$ for $j = 1, 2, i = 1, \dots, n$. Suppose that $\psi_{1i}(\partial \alpha_i) = \psi_{2i}(\partial \alpha_i)$ and $\psi_{1i}(\alpha_i)$ and $\psi_{2i}(\alpha_i)$ are ambient isotopic in B^3 relative to ∂B^3 . Then we say that a local move $(\bigcup_{i=1}^n \psi_{1i}(\alpha_i), \bigcup_{i=1}^n \psi_{2i}(\alpha_i))$ is a *parallel of (T_1, T_2) with weight $(l_1, \dots, l_n)$* .




Fig. 3.1

Proposition 3.2. *Let G be a connected, planar, prime and trivalent graph. Let $f_i : G \rightarrow S^3$ ($i = 1, 2$) be embeddings and S_i the canonical disk/band surface for f_i ($i = 1, 2$). If f_1 and f_2 are C_k -equivalent, then ∂S_1 and ∂S_2 are C_k -equivalent.*

Let T_1 and T_2 be tangles. We say that T_2 is obtained from T_1 by a local move (U_1, U_2) if there is an orientation preserving embedding $h : B^3 \rightarrow \text{int} B^3$ such that $S_i \cap h(B^3) = h(U_i)$

for $i = 1, 2$ and $T_1 - h(B^3) = T_2 - h(B^3)$. Two tangles T_1 and T_2 are C_k -equivalent if T_2 is obtained from T_1 by a finite sequence of C_k -moves and ambient isotopies relative ∂B^3 .

Lemma 3.3. (cf. [3, Claim on p. 26]) *Let (T_1, T_2) be a parallel of a C_k -move. Then T_1 and T_2 are C_k -equivalent.*

Proof. Let (U_1, U_2) be a C_k -move, $u_{11}, \dots, u_{1k+1}$ and $u_{21}, \dots, u_{2k+1}$ the components of U_1 and U_2 respectively. Suppose that (T_1, T_2) is a parallel of (U_1, U_2) with weight $(l_1, \dots, l_{k+1})$. We give a proof by induction on $l_1 \times \dots \times l_{k+1}$. In the case that $l_1 \times \dots \times l_{k+1} = 1$, it obviously holds. Suppose $l_1 \times \dots \times l_{k+1} \geq 2$. We may suppose $l_1 \geq 2$. By Lemma 2.1 in [17], we may assume that the C_k -move (U_1, U_2) is as illustrated in Fig. 3.2, i.e., the arcs except for u_{i1} are contained in the shaded part in U_i ($i = 1, 2$). It is not hard to see that T_2 is obtained from T_1 by l_1 local moves that are parallels of (U_1, U_2) with weight $(1, l_2, \dots, l_{k+1})$. This completes the proof. $\square$




Fig. 3.2

Proof of Proposition 3.2. In the case $k = 1$, it clearly holds. We consider the case $k \geq 2$. It is sufficient to consider the case that f_2 is obtained from f_1 by a single C_k -move. Suppose that f_2 is obtained from f_1 by a C_k -move. Then there is an embedding $h : B^3 \rightarrow S^3$ such that $(h^{-1}(f_1(\theta)), h^{-1}(f_2(\theta)))$ is a C_k -move and $f_1(\theta) - h(B^3) = f_2(\theta) - h(B^3)$ together with the labels. We may suppose that $(h^{-1}(\partial S_1), h^{-1}(\partial S_2))$ is a parallel of the C_k -move $(h^{-1}(f_1(\theta)), h^{-1}(f_2(\theta)))$ with weight $(2, \dots, 2)$. Then we have a new disk/band surface S'_2 for f_2 from S_1 by the local move $(h^{-1}(\partial S_1), h^{-1}(\partial S_2))$. Since C_k -move ($n \geq 2$) does not change the linking number, by Theorem 3.1, we have $\langle f_1(c_i), f_1(c_j) \rangle_{S_1} = \langle f_2(c_i), f_2(c_j) \rangle_{S_2}$, and since the local move $(h^{-1}(\partial S_1), h^{-1}(\partial S_2))$ does not change the Seifert linking form of a disk/band surface, we have $\langle f_1(c_i), f_1(c_j) \rangle_{S_1} = \langle f_2(c_i), f_2(c_j) \rangle_{S'_2}$. So we have $\langle f_2(c_i), f_2(c_j) \rangle_{S_2} = \langle f_2(c_i), f_2(c_j) \rangle_{S'_2}$. By Theorem 3.1, S_2 is ambient isotopic to S'_2 . Thus ∂S_2 is obtained from ∂S_1 by a parallel of a C_k -move. Lemma 3.3 completes the proof. $\square$

Let G be a connected, planar, prime and trivalent graph and $\Gamma(G)$ the set of spatial graph types. Let v be an invariant of ordered, oriented links that takes values in an abelian group A . Then we define a map $s : \Gamma(G) \rightarrow A$ as $s(f) = v(\partial S)$, where S is the canonical disk/band surface for f . By Theorem 3.1, s is an invariant of f . We call s the *invariant induced from v* .

Theorem 3.4. *Let G be a connected, planar, prime and trivalent graph and $\Gamma(G)$ the set of spatial graph types. Let v be a Vassiliev invariant of type $(k_1, \dots, k_l)$ for ordered,*

oriented links. Then the invariant for $\Gamma(G)$ induced from v is a Vassiliev invariant of type $(k_1, \dots, k_l)$.

In Theorem 3.4, the case of that a graph is the θ -curve and $k_1 = \dots = k_l = 1$ is given by Stanford [11].

By the arguments similar to that in the proof of Lemma 1.4 in [11], we have

Lemma 3.5. *Let v be a Vassiliev invariant of type $(k_1, \dots, k_l)$ for ordered, oriented links and s the invariant induced from v . Let $\{f_P | P \subset \{1, \dots, l\}\}$ be a singular spatial graph of type $(k_1, \dots, k_l)$. Let L_P be the ordered, oriented link that is the boundary of the canonical disk/band surface for f_P ($P \subset \{1, \dots, l\}$). Suppose there are mutually disjoint embeddings $h_{ij} : B^3 \rightarrow S^3$ ($i = 1, \dots, l$, $j = 1, \dots, n_i$) such that*

- (1) $L_\emptyset - \bigcup_{i,j} h_{ij}(B^3) = L_P - \bigcup_{i,j} h_{ij}(B^3)$ together with orientations and labels of the components for any subset $P \subset \{1, \dots, l\}$,
- (2) $(h_{ij}^{-1}(L_\emptyset), h_{ij}^{-1}(L_{\{1, \dots, l\}}))$ is a C_{k_i} -move ($i = 1, \dots, l$, $j = 1, \dots, n_i$), and
- (3) $L_P \cap h_{ij}(B^3) = \begin{cases} L_{\{1, \dots, l\}} \cap h_{ij}(B^3) & \text{if } i \in P, \\ L_\emptyset \cap h_{ij}(B^3) & \text{otherwise.} \end{cases}$

Then we have

$$\sum_{P \subset \{1, \dots, l\}} (-1)^{|P|} s(f_P) = \sum_{P \subset \{1, \dots, l\}} (-1)^{|P|} v(L_P) = 0. \square$$

The following lemma follows directly from the proof of Theorem 1 in [10].

Lemma 3.6. *Let G be a connected, planar, prime and trivalent graph, $f_0 : G \rightarrow \mathbb{R}^2$ an embedding, and S_0 the regular neighborhood of $f_0(G)$ in $\mathbb{R}^2$. Let S be a disk/band surface for an embedding f . Suppose that S is the image of an embedding of S_0 that is an extension of f . Then the canonical disk/band surface for f is obtained from S by a finite sequence of the moves as illustrated in Fig. 3.3. $\square$*




Fig. 3.3

In the definition of band sum in Section 2, by replacing $f_i(\theta)$ with T_i ($i = 1, 2$), we can define that T_2 is a *band sum* of T_1 and link models $(\alpha_1, \beta_1), \dots, (\alpha_l, \beta_l)$. By the arguments similar to that in Proof of Lemma 3.6 [17], we have the following lemma.

Lemma 3.7. *Two tangles T_1 and T_2 are C_k -equivalent if and only if there are tangless T'_1 and T'_2 such that T'_i is ambient isotopic to T_i ($i = 1, 2$) relative ∂B^3 and T'_2 is a band sum of T'_1 and some C_k -link models. $\square$*

Lemma 3.8. *Let T_1 and T_2 be tangles. If T_2 is obtained from T_1 by a parallel of a C_k -move, then there are tangles T'_1, T'_2 and mutually disjoint, orientation preserving embeddings $h_i : B^3 \rightarrow \text{int} B^3$ ($i = 1, \dots, n$) such that*

- (1) T'_j is ambient isotopic to T_j ($j = 1, 2$) relative ∂B^3 ,

- (2) $T'_1 - \bigcup_i h_i(B^3) = T'_2 - \bigcup_i h_i(B^3)$, and
(3) $(h_i^{-1}(T'_1), h_i^{-1}(T'_2))$ is a C_k -move ($i = 1, \dots, n$).

Proof. By Lemma 3.3, T_1 and T_2 are C_k -equivalent. By Lemma 3.7, there are tangles T'_1 and T'_2 such that T'_j is ambient isotopic to T_j ($j = 1, 2$) relative ∂B^3 and that T'_2 is a band sum of T'_1 and some C_k -link models $(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)$. Since $(\alpha_i, \hat{\beta}_i)$ ($i = 1, \dots, n$) are C_k -moves, we have the conclusion. $\square$

Proof of Theorem 3.4. Since $\{f_P | P \subset \{1, \dots, l\}\}$ is a singular spatial graph of type $(k_1, \dots, k_l)$, by the definition, there are mutually disjoint, orientation preserving embeddings $h_i : B^3 \rightarrow S^3$ ($i = 1, \dots, l$) such that

- (1) $f_\emptyset(G) - \bigcup_i h_i(B^3) = f_P(G) - \bigcup_i h_i(B^3)$ together with the labels for any subset $P \subset \{1, \dots, l\}$,
(2) $(h_i^{-1}(f_\emptyset(G)), h_i^{-1}(f_{\{1, \dots, l\}}(G)))$ is a C_{k_i} -move ($i = 1, \dots, l$), and
(3) $f_P(G) \cap h_i(B^3) = \begin{cases} f_{\{1, \dots, l\}}(G) \cap h_i(B^3) & \text{if } i \in P, \\ f_\emptyset(G) \cap h_i(B^3) & \text{otherwise.} \end{cases}$

Let $S_\emptyset$ be the canonical disk/band surface for $f_\emptyset$. By considering the intersections $S_\emptyset \cap h_i(B^3)$ ($i = 1, \dots, l$), we find disk/band surfaces S_P for f_P ($P \subset \{1, \dots, l\}$) such that

- (1) $S_\emptyset - \bigcup_i h_i(B^3) = S_P - \bigcup_i h_i(B^3)$
(2) $(h_i^{-1}(\partial S_\emptyset), h_i^{-1}(\partial S_{\{1, \dots, l\}}))$ is a parallel of a C_{k_i} -move ($i = 1, \dots, l$), and
(3) $S_P \cap h_i(B^3) = \begin{cases} S_{\{1, \dots, l\}} \cap h_i(B^3) & \text{if } i \in P, \\ S_\emptyset \cap h_i(B^3) & \text{otherwise.} \end{cases}$

By the proof of Proposition 3.2, if $k_i \geq 2$ for any $i \in P$, then S_P is the canonical disk/band surface. Set $h_i = h_{i1}$ ($i = 1, \dots, l$). By Lemma 3.6, there are mutually disjoint, orientation preserving embeddings $h_{ij} : B^3 \rightarrow S^3$ ($i = 1, \dots, l$, $j = 1, \dots, n_i$), where $n_i = 1$ if $k_i \geq 2$, and the canonical disk/band surfaces S'_P for f_P such that

- (1) $S'_\emptyset - \bigcup_{i,j} h_{ij}(B^3) = S'_P - \bigcup_{i,j} h_{ij}(B^3)$,
(2) $(h_{ij}^{-1}(\partial S'_\emptyset), h_{ij}^{-1}(\partial S'_{\{1, \dots, l\}}))$ is a parallel of C_{k_i} -move ($i = 1, \dots, l$, $j = 1, \dots, n_i$), and
(3) $S'_P \cap h_{ij}(B^3) = \begin{cases} S'_{\{1, \dots, l\}} \cap h_{ij}(B^3) & \text{if } i \in P, \\ S'_\emptyset \cap h_{ij}(B^3) & \text{otherwise.} \end{cases}$

By combining this, Lemmas 3.8 and 3.5, we have the conclusion. $\square$

Let G be a connected, planar, prime and trivalent graph and $E(G) = \{e_1, \dots, e_n\}$ the set of edges of G . Let S_f be the canonical disk/band surface for a spatial embedding f of G , and let $S_f(x_1, \dots, x_n; y_1, \dots, y_n)$ ($x_i \in \mathbb{Z}, y_j \in \{-1, 0, 1\}$) be a surface obtained from S_f as illustrated in Fig. 3.4. We note that $S_f(x_1, \dots, x_n; y_1, \dots, y_n)$ depends only on S_f and the integers $x_1, \dots, x_n, y_1, \dots, y_n$. This means $S_f(x_1, \dots, x_n; y_1, \dots, y_n)$ is the unique surface for f . Let v be an invariant of ordered, oriented links that takes values in an abelian group A . Then we can define an invariant $s_{(x_1, \dots, x_n; y_1, \dots, y_n)} : \Gamma(G) \rightarrow A$ as $s_{(x_1, \dots, x_n; y_1, \dots, y_n)}(f) = v(\partial S_f(x_1, \dots, x_n; y_1, \dots, y_n))$. We call $s_{(x_1, \dots, x_n; y_1, \dots, y_n)}$ the *invariant induced from v with respect to $x_1, \dots, x_n, y_1, \dots, y_n$* .




Fig. 3.4

By the arguments similar to that in the proofs of Proposition 3.2 and Theorem 3.4, we have the following theorem.

Theorem 3.9 *Let G be a connected, planar, prime and trivalent graph and $E(G) = \{e_1, \dots, e_n\}$ the set of edges of G . Then the followings hold.*

(1) *Let f_1 and f_2 be spatial graphs and $S_i(x_1, \dots, x_n; y_1, \dots, y_n)$ the surface obtained from the canonical disk/band surface for f_i ($i = 1, 2$). If f_1 and f_2 are C_k -equivalent, then $\partial S_1(x_1, \dots, x_n; y_1, \dots, y_n)$ and $\partial S_2(x_1, \dots, x_n; y_1, \dots, y_n)$ are C_k -equivalent.*

(2) *Let v be a Vassiliev invariant of type $(k_1, \dots, k_l)$ for ordered, oriented links. Then the invariant $s_{(x_1, \dots, x_n; y_1, \dots, y_n)}$ for spatial embeddings of G induced from v is a Vassiliev invariant of type $(k_1, \dots, k_l)$. $\square$*

Proof of Theorem 1.8. Suppose that G_k is abelian. Let f_1 and f_2 be spatial θ -curves as illustrated in Fig. 3.5. Since G_k is abelian, $g = f_1 \# f_1 \# f_2$ and $h = f_1 \# f_2 \# f_1$ are C_k -equivalent. Then, by Theorem 1.1(1), $g - h \in \mathcal{V}(\underbrace{1, \dots, 1}_k)$. This means that g and h cannot

be distinguished by any Vassiliev invariant of order $\leq k - 1$. Let $S_g(1, 1, -1; 0, 0, 1)$ and $S_h(1, 1, -1; 0, 0, 1)$ be the surfaces obtained from the canonical disk/band surfaces for g and h respectively. We note that $\partial S_g(1, 1, -1; 0, 0, 1)$ and $\partial S_h(1, 1, -1; 0, 0, 1)$ contain pretzel knots $K_g = P(3, 3, -3, -2)$ and $K_h = P(3, -3, 3, -2)$ respectively, see Fig. 3.6. Let v be a Vassiliev invariant for oriented link of order $\leq k - 1$. By combining Theorem 3.9(1) and the fact that a C_k -moves preserves Vassiliev invariants of order $\leq k - 1$ [3] (or simply by Theorem 3.9(2)), we have $v(\partial S_g(1, 1, -1; 0, 0, 1)) = v(\partial S_h(1, 1, -1; 0, 0, 1))$. Hence $v(P(3, 3, -3, -2)) = v(P(3, -3, 3, -2))$. Let $Q_K(q)$ be the quantum invariant of a knot K corresponding to the representation of the partition $(2, 1)$ of the quantum enveloping algebra $\mathcal{U}(sl_4)$. Then, using the computer software ‘K2K’ by M. Ochiai and N. Imafuji [9], we have

$$\begin{aligned} \frac{Q_{K_g}(q) - Q_{K_h}(q)}{Q_O(q)} &= q^8(-1+q)^{11}(1+q)^{11}(1+q^2)^3(1-q+q^2)^3(1+q+q^2)^3 \\ &\quad \times (1+q^4)^2(1-q^2+q^4)(1-q+q^2-q^3+q^4-q^5+q^6)^2 \\ &\quad \times (1+q+q^2+q^3+q^4+q^5+q^6)^2(1+q^8) \\ &\quad \times (1-q^2+q^4-q^6+q^8-q^{10}+q^{12}), \end{aligned}$$

where O is a trivial knot. Since this is divisible by $(1-q)^{11}$ and is not divisible by

$(1 - q)^{12}$, these pretzel knots can be distinguished by a Vassiliev invariant of order ≤ 11 [1]. Hence we have $k < 12$. This completes the proof. $\square$




Fig. 3.5




Fig. 3.6

Acknowledgement

The author would like to thank Professor Jun Murakami, Professor Józef Przytycki and Professor Paweł Traczyk for their valuable advice on distinguishing two mutant knots by Vassiliev invariants. Their advice was useful for proving Theorem 1.8

References

1. J. Birman and X.-S. Lin: Knot polynomials and Vassiliev's invariants, *Invent. Math.*, 111, 225-270, 1993.
2. K. Habiro: Aru musubime no kyokusyo sousa no zoku ni tuite (in Japanese), Master thesis in Tokyo University, 1994.
3. K. Habiro: Claspers and finite type invariants of links, *Geom. Topol.*, 4, 1-83, 2000.
<http://www.maths.warwick.ac.uk/gt/GTVol4/paper1.abs.html>

4. T. Kanenobu: Vassiliev-type invariants of a theta-curve, *J. Knot Theory and Its Ramifications*, 6, 455-477, 1997.
5. L. Kauffman, J. Simon, K. Wolcott and P. Zhao: Invariants of theta-curves and other graphs in 3-space, *Topology Appl.* 49, 193-216, 1993.
6. A. Koike: Finite-type invariants of embeddings of a theta-curve up to 4, *Yokohama Math. J.*, 47, 245-252, 1999.
7. B. Mellor: Finite-type link homotopy invariants II: Milnor's $\bar{\mu}$ invariants, preprint.
math.GT/9812119
8. T. Motohashi and K. Taniyama: Delta unknotting operation and vertex homotopy of graphs in $\mathbb{R}^3$, Proceedings of Knots 96, (S. Suzuki ed.), World Sci. Publ. Co., 185-200, 1997.
9. M. Ochiai and N. Imafuji: KNOT2000(K2K)
ftp://ftp.ics.nara-wu.ac.jp/pub/ochiai
10. T. Soma, H. Sugai and A. Yasuhara: Disk/band surfaces of spatial graphs, *Tokyo J. Math.*, 20, 1-11, 1997.
11. T. Stanford: The functoriality of Vassiliev-type invariants of links, braids, and knotted graphs. Random knotting and linking (Vancouver, BC, 1993). *J. Knot Theory Ramifications* 3, 247-262, 1994.
12. T. Stanford: Finite-type invariants of knots, links and graphs, *Topology*, 35, 1027-1050, 1996.
13. T. Stanford: Braid commutators and delta finite-type invariants, preprint.
math.GT/9907071
14. S. Suzuki: A prime decomposition theorem for a graph in 3-sphere, *Topology and Computer Science*, (S. Suzuki ed.), Kinokuniya, 259-276, 1987.
15. K. Taniyama: Cobordism, homotopy and homology of graphs in $\mathbb{R}^3$, *Topology*, 33, 509-523, 1994.
16. K. Taniyama and A. Yasuhara: Local moves on spatial graphs and finite type invariants, preprint.
17. K. Taniyama and A. Yasuhara: Band description of knots and Vassiliev invariants, to appear in *Math. Proc. Cambridge Philos. Soc.*
math.GT/003021
18. K. Wolcott: The knotting of theta curves and other graphs in S^3 , *Geometry and Topology* (C. McCrory and T. Shifrin ed.), Marcel Dekker, 325-346, 1987.